

MA 103 CHAPTER 9: SECTION 9.4      PROPERTIES OF LOGARITHMS

In Section 9.2 we discovered that for  $b > 0$  and  $b \neq 1$ , and  $a > 0$ .

1.  $\log_b b = 1$
2.  $\log_b 1 = 0$
3.  $\log_b (a) = c$  is equivalent to  $b^c = a$
4. WRITE THE EXPONENTIAL FORM FOR
  - A.  $\log_4 64 = 3$
  - B.  $\log_k m = n$
5. WRITE THE LOGARITHMIC FORM FOR
  - A.  $3^2 = 9$
  - B.  $p^q = r$
6. POWER PROPERTY FOR LOGARITHMS
  - A. Does  $\log_3 3^2 = 2 \log_3 3$ ?
  - B. Does  $\log_7 7^4 = 4 \log_7 7$ ?
  - C. Use a calculator table to compare values for  $f(x) = \log(x^3)$  and  $g(x) = 3 \log(x)$ .
  - D. Can you guess the power property:  $\log(x^p) = \underline{\hspace{2cm}}$ ?

7. THE COMMON AND NATURAL LOGARITHMS

Note:  $\log_2 64 = 6$  while  $\log_4 64 = 3$ , obviously the answer to a logarithm of a number  $n$  will differ depending upon the base of the logarithm. Thus, you can't evaluate  $\log_b n$  using any other log base except for  $b$ .

Your calculator has two logarithmic buttons:

$\log$  for  $\log_{10}$  Referred to as the common log

$\ln$  for  $\log_e$  Referred to as the natural log

( $e$  is an irrational number which can be approximated by  $e = 2.7183$ ) (p. 239)

$\log(a) = c$  and  $10^c = a$  are equivalent

$\ln(a) = c$  and  $e^c = a$  are equivalent

Either of these logs can be used to solve exponential equations.

8. USING THE POWER PROPERTY TO SOLVE EXPONENTIAL EQUATIONS:

A.  $2^x = 24$

B.  $3^{7x+1} + 7 = 50$

C.  $5^x + 9 = 40 - 3(5^x)$

D.  $20 = -3 + 4(2^x)$

9. MISUSE OF THE POWER PROPERTY

It is true that  $\log_b (x^p) = p \log_b (x)$ .

Only one of the following statements is true, which one is it?

(1)  $\log_b [a(x^p)] = p \log_b (ax)$

(2)  $\log_b [(ax)^p] = p \log_b (ax)$

Let  $b = 10$ ,  $a = 10$ ,  $x = 10$  and  $p = 2$  and see if the result agrees with your guess.

10. SOLVING LOGARITHMIC EQUATIONS

A.  $\log_8 x = 1/3$

B.  $\log_5 (4x + 1) = 3$

C.  $\log_b (80) = 7$

D.  $\log_b (16) = 2$